

References

- Aurivillius, C. (1888-1892). *Nordens Fjärilar, handbok i Sveriges, Norges, Danmarks och Finlands Macrolepidoptera*. Aktiebolaget, Hiertas Bokförlag, Stockholm (p.143, 1891).
- Bjerkander, C. (1778). Om Hvitax-Masken. *K.svenska Vetensk.Akad. Handl.* **39**: 289-293.
- Dahlbohm, G. (1837). *Kort underrättelse om skandinaviska Insekters allmännare skada och nytta i hushållningen*. Fr. Berlings Forlag, Lund.
- Esper, E.J.C. (1786-1805). *Die Schmetterlinge in Abbildungen nach der Natur mit Beschreibungen*, **4**. Die Eulenphalenen. Wolfgang Walthers, Erlangen.
- Gustafsson, B. (ed.) (1987). *Catalogus Lepidopterorum Suecica*. Entomologiska Foreningen, Naturhistoriska Riksmuseet, Stockholm.
- Heinicke, W. (1960). Revision der Gattung *Apamea* Ochs, 1816 (Lep.: Noctuidae) 2. Teil II Die Formen von *Mesapamea secalis* L. *D.ent. Z. N.F.* **7**: 166-176.
- Horn, W. (1926). Über den Verbleib der entomologischen Sammlungen der Welt. *Supplta.ent.* **12**: 1-120.
- Kaaber, S. & Skule, B. (1985). Noctuidae. In Katalog over de danske Sommerfugle: 103-114. *Ent. Meddr.* **52** (2-3): 1-163.
- Linnaeus, C. (1758). *Systema Naturae*. ed. 10. Laurentius Sylvius, Stockholm.
- (1761). *Fauna Svecica*, editio altera. Laurentius Sylvius, Stockholm.
- (1767). *Systema Naturae*. ed. 12. Laurentius Sylvius, Stockholm.
- Remm, H. (1983). New species of Noctuidae (Lepidoptera) from the USSR. *Ent. Obazr.* **62**: 591-600.
- Rezbanyai-Reser, L. (1985). *Mesapamea*-Studien II. *Mesapamea remmi* sp.n. aus der Schweiz, sowie Beiträge zur Kenntnis der westpalaearktischen Arten der Gattung *Mesapamea* Heinicke 1959. (Lep.: Noctuidae). *Ent. Ber. Luzern.* **14**: 127-148.
- Robinson, G.S. & Nielsen, E.S. (1983). The microlepidoptera described by Linnaeus and Clerck. *Syst. Ent.* **8**: 191-242.
- Rolander, D. (1752). Hvit-ax-Masken. *K.svenska Vetensk.Akad. Handl.* **13**: 62-66.
- Schoyen, M.W. (1879). *Pyralis secalis* L. *Stettin. ent. Ztg.* **40**: 389-396.
- Staudinger, O. & Rebel, H. (1901). *Catalog der Lepidopteren des palaearktischen Faunengebietes*. R. Friedländer, Berlin.
- Tutt, J.W. (1891). *The British noctuae and their varieties*. **1**. Warne, London.

A NOTE ON TINAGMA BALTEOLELLA F.v.R. (LEP.: DOUGLASIIDAE) — In the autumn of 1987 I collected a number of dead stems of viper's bugloss (*Echium vulgare*) from the shingles at Dungeness, Kent, in order to breed and photograph the common, stem-feeding species *Tinagma ocnerosomella* Staint.

The stems were hung outside, exposed to the elements, all winter and brought inside late April 1988. On 9th May the first of a considerable number of *T. balteolella* emerged. This species was first added to the British list in 1976 (Agassiz, *Ent. Gaz.* **26**: 291-293) and, as far as I am aware, has only been noted from the coastal sand-dunes of east Kent, from where it was first recorded. Dungeness is geographically reasonably close to the original locality, but ecologically very different.

Perhaps it is worth looking further afield for this easily overlooked species? PAUL SOKOLOFF, 4 Steep Close, Orpington, Kent.